

Helium Clustering in Neutron-Rich Be Isotopes

N.I. Ashwood¹, M. Freer¹, S. Ahmed¹, J.C. Angélique², V. Bouchat³, W.N. Catford^{4,2}, N.M. Clarke¹, N. Curtis¹, O. Dorvaux⁵, B.R. Fulton⁶, F. Hanappe³, Y. Kerckx³, M. Labiche⁷, J.L. Lecovey^{2†}, R.C. Lemmon⁸, F.M. Marqués², T. Materna³, A. Ninane⁹, G. Normand², N.A. Orr², S. Pain⁴, N. Soić^{1‡}, L. Stuttgé⁵, C. Timis^{2*}, A. Unsharova¹⁰, J.S. Winfield¹¹ and V.A. Ziman¹

¹*School of Physics and Astronomy, University of Birmingham, Edgbaston, Birmingham, B15 2TT, United Kingdom.*

²*Laboratoire de Physique Corpusculaire, ISMRA and Université de Caen, IN2P3-CNRS, 14050 Caen Cedex, France.*

³*Université Libre de Bruxelles, CP 226, B-1050 Bruxelles, Belgium.*

⁴*School of Electronics and Physical Sciences, University of Surrey, Guildford, Surrey, GU2 7XH, United Kingdom.*

⁵*Institut de Recherches Subatomique, IN2P3-CNRS/Université Louis Pasteur, BP 28, 67037 Strasbourg Cedex, France.*

⁶*Department of Physics, University of York, Heslington, York, YO10 5DD, United Kingdom.*

⁷*Department of Electronic Engineering and Physics, University of Paisley, Paisley, PA1 2BE, United Kingdom.*

⁸*CLRC Daresbury Laboratory, Daresbury, Warrington, Cheshire, WA4 4AD, United Kingdom.*

⁹*Institut de Physique, Université Catholique de Louvain, Louvain-la-Neuve, Belgium.*

¹⁰*Joint Institute for Nuclear Research, 141980, Dubna, Moscow Region, Russia.*

¹¹*Instituto Nazionale di Fisica Nucleare, Laboratori Nazionali del Sud, 44 - 95123 Catania, Italy.*

(Received February 8, 2008)

Measurements of the helium-cluster breakup and neutron removal cross sections for neutron-rich Be isotopes ^{10–12,14}Be are presented. These have been studied in the 30 to 42 MeV/u energy range where reaction measurements are proposed to be sensitive to the cluster content of the ground-state wavefunction. These measurements provide a comprehensive survey of the decay processes of the Be isotopes by which the valence neutrons are removed revealing the underlying α - α core-cluster structure. The measurements indicate that clustering in the Be isotopes remains important up to the drip-line nucleus ¹⁴Be and that the dominant helium-cluster structure in the neutron-rich Be isotopes corresponds to α - Xn - α .

PACS number(s): 25.60.Dz, 25.70.-z, 25.70.Mn, 27.20.+n

The structure of nuclear matter at the limits of stability has always been a source of fascination. These limits are bounded by the extremes of excitation energy or spin, on one hand, and by the extremes of either total number of nucleons or ratios of protons to neutrons (isospin) on the other. Most notably, radioactive beam facilities provide access to the limit of neutron excess. Here, structural properties are known to differ from normal nuclear matter, as they are strongly influenced by the weak binding of the valence neutrons. The classic example is the halo, where weakly bound neutrons exist as a diffuse cloud around a compact core [1]. In this case there is a decoupling of the valence neutrons from a core, or cluster, of normal nuclear matter.

Recently, it has been speculated that such clustering may be a general feature of drip-line nuclei caused by the tendency of the nucleus to distribute the neutron excess among the core nucleons [2]. In order to maximize the overlap of the excess neutrons with protons, the core may either deform enhancing the surface area, or optimally, form clusters. Thus, the ground states of neutron drip-

line nuclei may resemble clusters embedded in a cloud of weakly bound neutrons.

Certain nuclei already possess a predisposition towards clustering. For example, ⁸Be clearly demonstrates a strong α - α cluster character [3,4]. Recent experimental and theoretical developments indicate that this underlying cluster structure has a strong influence on the neutron-rich Be isotopes. In particular, the two-centred nature of the potential in which the valence neutrons reside may produce molecular type characteristics in these nuclei [5,6]. Calculations using the antisymmetrized molecular dynamics (AMD) framework suggest such features can occur in the Be isotopes [7,8]. The same model predicts that the normally compact boron isotopes become highly clustered at the drip-line [9], in line with the ideas concerning the optimization of the proton valence-neutron overlap [2].

Direct access to such structural properties of the ground states of very neutron-rich nuclei is extremely challenging due to the fact that the ^xHe cluster breakup thresholds increase significantly towards the drip-line. However, the AMD model has been used to explore how the fragmentation of beams of boron isotopes may be used to extract a signature for the clustering in the ground state [10]. This study revealed that such clustering may be probed via fragmentation if energies in the region of 30 MeV/nucleon were used. However, studies of the most neutron-rich boron isotope (¹⁹B), as suggested in [10], is beyond the capability of present day experimental facilities. It is, however, possible to study the beryllium nuclei all the way from stability to the drip-lines. Here we present comprehensive measurements of ^xHe-cluster, which were explored theoretically for the boron isotopes, and neutron breakup cross sections for the beryllium isotopes A=10, 11, 12 and 14. These data should provide a significant test of our understanding of

the structure of nuclei at the neutron drip-line.

The measurements were performed at the GANIL accelerator facility with the production of $^{10,11,12,14}\text{Be}$ secondary beams via the fragmentation of ^{13}C and ^{18}O primary beams. The reaction products were mass, charge and momentum analyzed using the LISE3 spectrometer producing beams of purity $\geq 95\%$ for the lighter isotopes and $\simeq 20\%$ for the heaviest and count rates of the order of 10^4 pps for masses 10, 11 and 12 (limited by the count rate capacity of the detection system) and ~ 50 pps for ^{14}Be . Identification of the beam of the beam particle was achieved using time-of-flight through the LISE separator as measured by a PPAC at the entrance to the reaction chamber. The energies of the 4 beams were 30.9, 41.7, 41.8 and 34.4 MeV/u in order of increasing mass.

The beam was tracked onto 20 to 275 mg/cm² carbon targets using two drift chambers, providing a measurement of the incident position and angle of the projectile. The beam and reaction products then entered a zero-degree telescope formed from two 500 μm thick, 16-strip position-sensitive silicon detectors placed 15 cm from the target. These two detectors were arranged with orthogonal strips, providing a measurement of the incident ions to ≤ 1 mm in both the x and y directions (the z -coordinate being the beam direction). Behind the strip detectors was a close packed array of 16 1.5 cm thick, 2.5×2.5 cm², CsI scintillators. These were placed at 30 cm from the target so as to cover the same solid angle as the strip detectors as measured from the target. Calibration of the energy response of the silicon and CsI detectors was achieved using α -sources and a mixed beam of light-ions of known energy. Neutrons produced in reactions of the beam were detected using an array of ~ 100 neutron detectors (D $\acute{\text{e}}$ MoN) arranged in a similar manner to that shown in [11]. These detectors gave a single-neutron detection efficiency of $\sim 15\%$ (as in [13]). Monte Carlo simulations of the response of the charged particle detection system indicates that the efficiency for the detection of the breakup of the Be isotopes into two ^xHe nuclei was 40 to 50 % and almost independent of excitation energy of the decaying system. The measured relative energy of the decay products ($E_{\text{rel}} = \mu v_{\text{rel}}^2/2$) thus allowed the excitation energy to be deduced ($E_x = E_{\text{rel}} + E_{\text{thresh}}$).

A knowledge of the breakup yields, the target thickness, detection efficiency and the integrated beam exposure allowed the cross sections for the various breakup processes to be calculated. In addition, as noted above, it was possible to deduce the excitation energy of the decaying system. As an example, Fig. 1 shows the $^4\text{He}+^4\text{He}$ invariant mass spectrum (E_x) for the $^{12,14}\text{Be}$ beams. It is clear that a large fraction of the events involving the dissociation of ^{14}Be (^{12}Be) proceed via 6 (4) neutron emission to form ^8Be , i.e. the decay process does not always follow from the decay into neutron-rich helium isotopes followed by neutron emission, i.e. unbound excited states

of ^xHe . The evidence for this is the observation of peaks in the ^8Be excitation energy spectrum at energies which correspond to the decay of the ground-state (92 keV) and the first excited state (3.04 MeV, 2^+). There is one further feature in this spectrum, at ~ 600 keV, which does not correspond to a resonance in ^8Be but is produced by the decay of the 2.43 MeV ($5/2^-$) state in ^9Be to the tail of the 2^+ state in ^8Be . The presence of this feature suggests a fraction of the yield proceeds via sequential 6n emission to ^8Be . Similar features are also present in the $^4\text{He}+^4\text{He}$ decay spectra measured with the other Be projectiles.

The mass identification provided by the zero degree telescope allows the resolution of all the He isotopes and thus permits all stages of the decay process to be reconstructed. For example, in the case of ^{14}Be it is possible to reconstruct the decay into $^6\text{He}+^8\text{He}$, or following the emission of neutrons from the projectile, the decay of $^{12}\text{Be}^*$ into $^6\text{He}+^6\text{He}$ (or $^4\text{He}+^8\text{He}$), or the decay of $^{10}\text{Be}^*$ to $^6\text{He}+^4\text{He}$, or finally, as discussed above, ^8Be to $^4\text{He}+^4\text{He}$. The cross sections for these processes are plotted in Fig. 2. These cross sections are also presented together with the $^x\text{Be}+1\text{n}$ (i.e. core fragment plus one neutron) cross sections in Table 1. The neutron cross sections have been calculated by selecting beam velocity neutrons ($E_n > 0.37 E_{\text{beam}}/A$). In the case of $^{11,12}\text{Be}$ these were calculated by integrating the neutron angular distributions with a Lorentzian single neutron line-shape, whilst subtracting the background from reactions within the CsI component of the telescopes, as deduced from measurements with no target [13]. In the case of $^{10,14}\text{Be}$ data no background measurements were performed and thus the cross sections provides upper limits only. For ^{14}Be , measurements of these cross sections, including the background subtraction, have already been made by Labiche *et al.* [12,13] at 35 MeV/u, and these are also shown in Table 1. Given an analysis of the no target yield from the ^{12}Be data (~ 20 -30% of events in 2n removal channel are from the target) the cross sections are consistent. Comparison of the cross sections with earlier studies show reasonable agreement. The 0.18 ± 0.11 b ^{11}Be 1n removal cross section compares with that extracted by Anne *et al.* [14] of 0.12 ± 0.02 b, and the 0.11 ± 0.04 b ^{12}Be 1n removal is in close agreement with that measured by Navin *et al.* [15], 0.045 ± 0.05 b, albeit at 78 MeV/u. Similar agreement is found with the helium breakup measurements reported in [16].

The measurement of the multi-neutron removal channels via the detection of the core plus one neutron presents the advantage that the detection efficiency does not become prohibitively small. Naively, the true cross section for the $A - Xn$ channel should be reduced by a factor X . However, the neutron angular distributions vary from decay step to decay step and thus the efficiency is not constant, and, moreover, the multiplicity of the emitted neutrons does not necessarily equal the num-

ber of missing neutrons, with the possibility of projectile neutrons interacting strongly with the target [13]. Indeed, the measured neutron multiplicities for $^{14}\text{Be} \rightarrow ^{12}\text{Be}$ and $^{14}\text{Be} \rightarrow ^{10}\text{Be}$ are found to be 1.63 ± 0.26 and 2.9 ± 0.8 [12,13]. Although the error bars are large these multiplicities indicate that there is a tendency for there to be slightly less neutrons in the final state than anticipated. Nevertheless, for the sake of the comparison with the measured helium breakup cross sections the $A-X\text{Be}+1n$ cross sections are plotted in Fig. 2 divided by X , thus indicating the strength with which the bound states of the $A-Xn$ nucleus are populated.

It is clear that neutron removal dominates even in the case of the nucleus ^{10}Be in which the neutron decay threshold (6.8 MeV) is almost commensurate with that for α -decay (7.4 MeV). In all cases the breakup of the projectile into helium fragments is several orders of magnitude smaller than the neutron removal cross sections. The helium breakup cross sections steadily increase as the neutron removal process feeds states above the cluster decay thresholds. At the point at which the systems arrive at ^8Be the cluster breakup and neutron removal processes coincide. For each step the ratio of yield in the $A-Xn$ and helium breakup gives an idea of the relative excitation probabilities to states above the neutron and cluster decay thresholds, and subsequent stages reveal the probability that the daughter $A-Xn$ system is formed in either bound states or states unbound to either neutron or helium decay. These decay systematics should allow detailed modeling of the decay process via which these nuclei lose all of their neutrons leaving the valence α - α cluster core.

Fig. 3 shows the mass dependence of the cluster decay cross sections. It is not possible to plot the data for ^{11}Be in this instance as the main decay channels $^5\text{He}+^6\text{He}$ $^4\text{He}+^7\text{He}$ lead to a $^4\text{He}+n+^6\text{He}$ final state which cannot be unambiguously distinguished from neutron removal followed by α -decay of ^{10}Be . Figs 3a and 3b show the cross sections for the helium cluster breakup of the projectile (first-chance breakup), i.e. $^{14}\text{Be} \rightarrow ^6\text{He}+^8\text{He}$, $^{12}\text{Be} \rightarrow ^6\text{He}+^6\text{He}$, $^{10}\text{Be} \rightarrow ^6\text{He}+^4\text{He}$, and the total He+He breakup cross section respectively. The first chance breakup cross section is essentially mass independent. These systematics indicate that the di-cluster content of the nuclei remains relatively constant from ^{10}Be through to ^{14}Be . However, the total breakup cross section for $A=10, 11, 12$ and 14 shows a strong minimum at $A=11$ increasing to maximum values for ^{10}Be and ^{14}Be . The ratio of first chance to total cluster breakup thus reveals a maximum at $A=12$.

Early calculations of the degree of clustering in the beryllium isotopes using the AMD framework indicated a decrease in clusterization from ^8Be to ^{12}Be with a small increase for ^{14}Be [8], with something like a factor of 2 change predicted in the clusterization between the ground states of ^{10}Be and ^{12}Be . However, most re-

cent studies using this framework now indicate low-lying states possess strong cluster symmetries [17]. If, as suggested by the AMD calculations for the reactions with the boron isotopes, the breakup process is strongly influenced by the overlap of the cluster content of the ground state and that of excited states above the decay threshold, then the present measurements indicate that there is no change in the degree of clustering in the mass range 10 to 14.

By far the most dominant He-cluster breakup channel is that corresponding to the removal of all of the valence neutrons leaving the α - α core (Fig 2). Table 2 shows the energy thresholds for $^8\text{Be}+Xn$ and first-chance He-cluster breakup. Remarkably, even though the thresholds for removing all valence neutrons are higher in energy, it is this channel which dominates, suggesting a strong influence of the α - Xn - α structure. Such a result cannot be explained by decay phase-space alone or the fact that with high numbers of valence neutrons there are many more partitions for the system to decay into. For example, in the case of ^{14}Be the decay threshold for $6n$ emission lies 4.2 MeV higher than that for cluster decay, and inelastic excitation probabilities would be expected to decrease with excitation energy. The ^{14}Be nucleus must emit one of six possible neutrons to resonant states above the ^{13}Be $5n$ decay threshold (10.3 MeV), rather than to lower energy states. In turn ^{13}Be must decay to resonances above the ^{12}Be $4n$ decay threshold. Thus, although there is a factor of 6 enhancement in the number of possible decay paths for the neutron emission, there are 5 intermediate steps in which the system may decay to states below the $\alpha+\alpha+6n$ threshold. Moreover, the phase space at each step is highly constrained.

It is possible that the excitation mechanisms leading to the first-chance He-cluster breakup and neutron removal are radically different. Certainly, direct elastic breakup and absorptive processes dominate in the first step over resonant breakup as they have been used extensively to determine the ground state structure of neutron-rich nuclei [1]. It is thus probable that a significant fraction of the $\alpha+\alpha+6n$ yield for ^{14}Be is produced via such processes.

The measurements presented here provide evidence for the existence of di-cluster structures in $^{10-12,14}\text{Be}$. Moreover, they appear to demonstrate that these nuclei possess a strong structural overlap with an α - Xn - α configuration. The comprehensive measurements of the neutron-removal and cluster breakup for the first time provide experimental data whereby the structure of the most neutron-rich Be isotopes can be modeled via their reactions.

The authors are grateful to the technical and operations staff of LPC and GANIL. This work was funded by the EPSRC (UK) and the IN2P3-CNRS (France). Additional support was provided by the Human Capital and Mobility Programme of the European Community (Con-

[†] Present address: NSCL, Michigan State University, Michigan 48824, U.S.A.

[‡] Present address: Rudjer Bošković Institute, Bijenička 54, HR-10000 Zagreb, Croatia.

* Present address: School of Electronics and Physical Sciences, University of Surrey, Surrey, GU2 7XH, U. K.

- [1] P.G. Hansen, A.S. Jensen, B. Jonson, *Annu. Rev. Nucl. Part. Sci.* **45**, 591 (1995).
- [2] H. Horiuchi, *Proceedings of the 7th International Conference on "Clustering Aspects of Nuclear Structure and Dynamics"*, World Scientific, 405, (2000).
- [3] B. Buck, H. Friedrich, and C. Wheatley, *Nucl. Phys.* **A275**, 246 (1977).
- [4] J. Hiura and R. Tamagaki, *Suppl. Prog. Theo. Phys.* **52**, 25 (1972).
- [5] W. von Oertzen, *Z. Phys.* **A357**, 355 (1997).
- [6] N. Itagaki and S. Okabe, *Phys. Rev. C* **61**, 044306 (2000).
- [7] Y. Kanada-En'yo, H. Horiuchi, and A. Doté, *Phys. Rev. C* **60**, 064304 (1999); A. Doté, H. Horiuchi, and Y. Kanada-En'yo, *Phys. Rev. C* **56**, 1844 (1997).
- [8] Y. Kanada-En'yo, H. Horiuchi, and A. Ono, *Phys. Rev. C* **52**, 628 (1995).
- [9] Y. Kanada-En'yo and H. Horiuchi, *Phys. Rev. C* **52**, 647 (1995).
- [10] H. Takemoto, H. Horiuchi, and A. Ono, *Phys. Rev. C* **63**, 034615 (2001).
- [11] F. M. Marqués *et al.*, *Phys. Rev. C* **65**, 044006 (2002).
- [12] M. Labiche, Ph.D thesis Laboratoire de Physique Corpusculaire, 1999, unpublished.
- [13] M. Labiche *et al.*, *Phys. Rev. Lett.* **86**, 600 (2001).
- [14] R. Anne *et al.*, *Nucl. Phys.* **A575**, 124 (1994).
- [15] A. Navin *et al.*, *Phys. Rev. Lett.* **85**, 266 (2000).
- [16] M. Freer *et al.*, *Phys. Rev. C* **63**, 034301 (2001).
- [17] Y. Kanada-En'yo and H. Horiuchi, *Phys. Rev. C*, accepted for publication.

	1n	2n	3n	4n	${}^6\text{He}+{}^8\text{He}$	${}^6\text{He}+{}^6\text{He}$	${}^8\text{He}+{}^4\text{He}$	${}^6\text{He}+{}^4\text{He}$	${}^4\text{He}+{}^4\text{He}$			Total He+He
									${}^8\text{Be}_{gs}$	${}^9\text{Be}(5/2^-)$	${}^8\text{Be}(2^+)$	
${}^{14}\text{Be}$		2248(82)	670(23)	799(26)	1.17(0.20)	1.33(0.22)	2.15(0.30)	10.04(0.65)	4.40(0.41)	5.40(0.44)	16.61(1.19)	41.09(1.54)
${}^{14}\text{Be}^\dagger$		750(10)		420(10)								
${}^{12}\text{Be}$	113(43)	237(54)	100(35)			0.23(0.02)	1.02(0.04)	3.04(0.05)	2.89(0.10)	4.25(0.10)	16.42(0.37)	27.86(0.40)
${}^{11}\text{Be}$	175(110)	22(12)						1.17(0.07)	3.59(0.14)	4.52(0.11)	13.12(0.31)	22.40(0.36)
${}^{10}\text{Be}$	29(4)							1.26(0.06)	7.97(0.40)	7.32(0.30)	21.6(1.1)	38.15(1.36)

	Neutron removal						He+He cluster decay
	1n	2n	3n	4n	5n	6n	
${}^{14}\text{Be}$	2.97	1.12	4.29	4.79	11.60	13.27	${}^6\text{He}+{}^8\text{He}$ 9.09
${}^{12}\text{Be}$	3.17	3.67	10.48	12.15			${}^4\text{He}+{}^8\text{He}$ 8.95
${}^{11}\text{Be}$	0.50	7.32	8.98				${}^5\text{He}+{}^6\text{He}$ 8.81
${}^{10}\text{Be}$	6.81	8.48					${}^4\text{He}+{}^6\text{He}$ 7.41

Figure and Table Captions

Table 1. Cross sections measured for He-cluster breakup and core plus one neutron in units of mb. Note in the case of ${}^{10,14}\text{Be}$ the neutron cross sections provide upper limits. The row marked ${}^{14}\text{Be}^\dagger$ are the background subtracted cross sections from [12,13].

Table 2. Energy thresholds for cluster and neutron decay in units of MeV.

Figure 1. The excitation energy spectrum corresponding to $\alpha+\alpha$ coincidences from the ${}^{12}\text{Be}$ beam. The peaks correspond to the decay of the ground-state (92 keV) and the first excited state (3.04 MeV, 2^+). The peak at ~ 600 keV is produced by the decay of the 2.43 MeV ($5/2^-$) state in ${}^9\text{Be}$ to the tail of the 2^+ state in ${}^8\text{Be}$. The inset is the excitation energy spectrum corresponding to $\alpha+\alpha$ coincidences from the ${}^{14}\text{Be}$ beam.

Figure 2. Normalized neutron-removal (open diamonds) and cluster breakup (filled circles) cross sections for the neutron-rich Be isotopes. For ${}^{14}\text{Be}$ the filled diamonds correspond to the measurements of [12,13]. Note, the points for the ${}^9\text{Be}$ breakup (open circles) are lower limits deduced from the 600 keV peak in the ${}^8\text{Be}$ spectrum in Figure 1. The vertical dotted lines indicate the mass of the projectile.

Figure 3. (a) The first-chance cluster breakup cross sections, i.e. ${}^{14}\text{Be} \rightarrow {}^6\text{He}+{}^8\text{He}$, ${}^{12}\text{Be} \rightarrow ({}^6\text{He}+{}^6\text{He}$ and ${}^4\text{He}+{}^8\text{He})$ and ${}^{10}\text{Be} \rightarrow {}^6\text{He}+{}^4\text{He}$, (b) the total He+He breakup cross sections.





